Abstract

Introduction

Earth and Venus have many similar features, yet their tectonic histories

are quite different. Like the Earth, Venus has a global rift system, which

has been cited as evidence of tectonic activity, despite the apparent lack

of Earth-style plate tectonics. Both systems are marked by large ridges,

usually with central grabens. On Earth, the topography of the rifts can be

modeled well by a cooling half-space and the spreading of two divergent

plates. The origin of the topographic signature on Venus, however, remains

circle arcs extending 1000s of kilometers. This chasma system measures

both have large regions of apparent upwelling - hotspots on Earth, and

regiones on Venus [Stofan and Smrekar, 2005]. These are marked by

broad topographic and geoid highs as well as evidence of volcanic activity.

We use topographic profiles to look for well-understood terrestrial analogs

to venusian features. Specifically, we cross-correlate average profiles for

terrestrial rifts (slow and fast, incipient and inactive) and hotspots (oceanic

and continental) with those for venusian chasmata and regiones, and draw

inferences as to the processes responsible for shaping Venus' surface.

determined for Earth [Jurdy and Stefanick, 1999]. Venus and Earth also

54,464 km [Parsons, 1981], which, when corrected for the smaller size of

enigmatic. Venus' rift zones (termed "chasmata") can be fit by four great

We use Principal Component Analysis (PCA) as an objective tool in analyzing, comparing, and contrasting topographic profiles of different/similar features from different locations and planets.

To do so, we take average profiles of a set of features and form a cross-correlation matrix, which is then diagonalized to determine its principal components. These components, not merely numbers, represent actual profile shapes that give a quantitative basis for comparing different sets of features. For example, PCA for terrestrial hotspots shows the main component as a generic dome shape. Secondary components show a more sinusoidal shape, related to the lithospheric loading response, and thus give information about the nature of the lithosphere setting of the various hotspots.

We examine a range of terrestrial spreading centers: fast, slow, ultra-slow, incipient, and extinct, and compare these to several chasmata on Venus (including Devana, Ganis, Juno, Parga, and Kuanja). For upwelling regions, we consider the oceanic Hawaii, Reunion, and Iceland hotspots and Yellowstone, a prototypical continental hotspot. Venus has approximately one dozen broad topographic and geoid highs called regiones. Our analysis includes Atla, Beta, and W. Eistla regiones. Atla and Beta are widely thought to be the most likely to be currently or

Analysis of terrestrial rifts suggests shows increasing uniformity of shape among rifts with increasing spreading rates. Venus' correlations of uniformity rank considerably lower than the terrestrial ones. Extrapolating the correlation/spreading rate suggests that Venus' chasmata, if analogous to terrestrial spreading centers, most resemble the ultra-slow spreading level (less than 12mm/yr) of the Arctic Gakkel ridge. PCA will provide an objective measurement of this correlation.

800-km profile lines used

Principal Component Analysis of Terrestrial and Venusian Topography

P.R. STODDARD, Department of Geology and Environmental Geosciences, Northern Illinois University, DeKalb, IL, USA D.M. Jurdy, Department of Earth and Planetary Sciences, Northwestern University, Evanston, IL, USA







Pac/Ant 0.678 0.295 0.028

Parga 0.423 0.189 0.388

Pac/Naz 0.715 0.269

Smith, D.E. and Zuber, M. T., 1996. The shape of Mars and the topographic signature of the hemispheric dichotomy. Science, 271, 184-188, DOI:10.1126/science.271.5246.184.

Lay, D. C., 2006, Linear Algebra and its Applications, New York, Pearson, Addison-Wesley, 499

Venus/Earth Rift Comparison **Terrestrial Rifts** Feature PC 1 PC 2 PC 3 PC 1 | PC 2 | Afr/NAm 0.663 0.229 0.108 E. Afr Rift 0.454 0.060 -0.486

Top 3 PC values, normalized so that absolute values sum to 1 (used as co-ordinates for ternary diagrams at left). Color coded to previous diagrams.

Using Magellan topography data for Venus, and ETOP05 for Earth, and techniques described more fully in Stoddard and Jurdy [2009], we compare profiles of venusian chasmata (Devana, Ganis, Parga, Juna, 'Lambda'') terrestrial spreading rifts: mid-ocean ridges, both fast (Pac/Naz, Coc/Pac, Pac/Ant) and slow (NAm/Eur, Afr/NAm, Afr/SAm); the ultra-slow, intermittent Arctic ridge; continental E. African Rift, and the extinct Labrador rift

First, we analyze this set of terrestrial spreading features with known spreading rates. Next, we compare a subset of Earth ridges (fast, intermediate, slow, ultra-slow and continental) with an equal number of Venus' chasmata: Ganis Chasma on Atla Regio and Devana Chasma, which extends from Beta Regio to Phoebe Regio [Price and Suppe, 1995], and Juno, Parga, and Lambda. On Venus, these rifts rank among Venus, nearly matches the 59,200-km total length of the spreading ridges | Items | the most recently active features [Basilevsky and Head, 2002] with profuse volcanism as documented by the nature of cratering.

> We cross-correlate the average profiles of each feature with each other to arrive at our correlation matrix, diagonalizing the matrix for the eigenvalues, or principal components (PCs). Each feature's profile can be sythesized by a weighted sum of the PC profiles.

- average profiles
- correlate with each other
- form correlation matrix
- diagonalize for PC analysis synthesize each feature

Ternary Diagrams for Principal-Component Analysis

We use ternary diagrams modified to allow plotting of variables with negative values to display relative strengths of the top three principal components (PC). The center triangle is the standard ternary, with all three variables (principal components) positive. Each of the adjacent three triangles has one negative variable. Each of the outermost three triangles has two negative variables. There were no features having three negative variables.

Covariance Matrix for Terrestrial Rifts

Top: Principal component profiles, in order of relative importance (largest eigenvalues at top). Bottom: Topographic profiles (black lines) of features analyzed in this study compared with profiles synthesized from PC profiles (blue lines). Left: Terrestrial rifts. Right: Venusian and terrestrial data set.

Ternary diagrams for terrestrial rifts (left) and selected terrestrial rifts with an equal number of venusian

Earth Ridges

chasmata (right).

Results

TERRESTRIAL SPREADING CENTERS

- PC decomposition fits very well:
- a. The top 3 PCs contain 98% of the topographic shape information
- b. The top 4 PCs contain 100%
- PC1 shows smooth simple uplift shape

Labrador -0.369 -0.136 -0.496

Pac/Ant 0.610 0.151 -0.239

NAm/Eur 0.595 0.164 -0.240

Pac/Naz 0.478 0.077 -0.445

- a. PC1 is positive for all continuously active spreading ridges, including the E. Afr. Rift
- b. PC1 is negative only for Labrador (extinct) and the Arctic (intermittent, ultra slow)
- PC2 reflects Arctic rift topography
- PC3 is dominated by a central trough
- a. PC3 is negative for all active mid-ocean ridges
- b. PC3 is positive for the E. African Rift
- Ternary diagram ("PC-space"):
- a. actively spreading ridges cluster
- b. E. Afr. Rift (continental), Labrador (extinct), and Arctic (ultra slow) are isolated in PC space

EARTH SPREADING CENTERS AND VENUS CHASMATA

- PC decomposition fits at a lower level; i.e., 88% for PCs 1-3; 95% for PCs 1-5
- PCs are less-well defined than for terrestrial-only case
- Earth's active spreading centers cluster
- Venus chasmata scatter; Devana and "Lambda" plot near E. African Rift
- PC1 is generally positive for Venus' features and Earth's active ridges
- PC3 to be negative for chasmata

entering transportation to the section of


Earth and Venus

Phoebe

285E

Conclusions

- Active mid-ocean ridges on Earth have similar and consistent topographic profile shapes; allows for a better fit of each profile using fewer PCs
- Venus chasmata and Earth's non-typical ridges display much more variability in topographic profile Since chasmata plot more like the Arctic and E. African rifts, we suggest that chasmata are ultra-slow spreading rifts in a thick lithosphere. A previous topographic study (REF) found that Atla and Beta regiones most closely resembles Yellowstone, another indication of a thick lithosphere for Venus



Average profiles for the 15 rifts on Earth (blue) and Venus

Average Profiles